




































































Willow Collection ProSpecieRara -- Overview

OVERVIEW regarding USAGE and CULTIVATION

| BEZEICHNUNG | | | VERWENDUNG | | | | | | | | | | | | KULTUR | | | | | |
|--|---|-----------|------------|---------------|---------|-----|-------|-----|-----|-----|-----|---------------------------------|-----|-----|--------|-----|-------|-------|-----|-----|
| Salix alba 'Capriasca' White Willow 'Capriasca' | M | Lv CH | --- | --- | | --- | ♀ XXX | --- | --- | ♂ | --- | --- | --- | --- | --- | --- | --- | 15-25 | ♂ | --- |
| Salix alba 'Soorwiesli' White Willow 'Soorwiesli' | M | Lv CH | --- | --- | | --- | ♀ XXX | --- | --- | ♂ | --- | --- | --- | --- | --- | --- | --- | 15-25 | ♂ | --- |
| Salix daphnoides 'Aglaiä' Violet Willow 'Aglaiä' | M | Cv EU | --- | --- | | --- | --- | --- | --- | ♂ | NP | Feb - - Mar 4 4 Apr 4 4 May - - | --- | --- | --- | --- | 10-15 | ♂ | --- | |
| Salix daphnoides 'Netta Statham' Violet Willow 'Netta Statham' | M | Cv GB | ♂ | --- XX X S | 1.5-2.1 | --- | --- | --- | --- | ♂ | NP | Feb - - Mar 4 4 Apr 4 4 May - - | --- | --- | --- | --- | 6-10 | ♂ | ♂ | |
| Salix nigricans 'Black Lady' Dark-leaved Willow 'Black Lady' | F | Lv CH | ♂ | --- X XXX S | 1.2-1.7 | ♀ | XXX | --- | --- | --- | --- | --- | --- | --- | --- | --- | 3-4 | --- | ♂ | |
| Salix nigricans 'Faucille' Dark-leaved Willow 'Faucille' | F | Cv | ♂ | --- X XXX S | 1.2-1.5 | ♀ | XX | --- | --- | --- | --- | --- | --- | --- | --- | --- | 3-4 | --- | ♂ | |
| Salix purpurea 'Carl Jensen' Purple Willow 'Carl Jensen' | M | Cv DK | ♂ | --- XX XXX | 1.6-1.9 | --- | --- | --- | --- | ♂ | --- | --- | --- | --- | --- | ? | 4-5 | --- | ♂ | |
| Salix purpurea 'Dark Dicks' Purple Willow 'Dark Dicks' | F | Cv GB | ♂ | --- --- XXX | 1.5-1.8 | ♀ | X | --- | --- | --- | --- | --- | --- | --- | --- | ? | 3-4 | --- | ♂ | |
| Salix purpurea 'Light Dicks' Purple Willow 'Light Dicks' | M | Cv GB | ♂ | --- X XXX | 1.5-1.8 | ♀ | X | --- | --- | --- | --- | --- | --- | --- | --- | ? | 3-4 | --- | ♂ | |
| Salix purpurea 'Uralensis' Ural Willow | F | Cv D/A | ♂ | --- XX XX | 1.5-2.5 | ♀ | XXX | --- | --- | --- | --- | --- | --- | --- | --- | ? | 3-4 | --- | ♂ | |
| Salix triandra 'Black German' Almond-leaved Willow 'Black German' | F | Cv GB | ♂ | XX --- --- SS | 1.8-2.5 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 5-7 | --- | ♂ | |
| Salix triandra 'Black Maul' Almond-leaved Willow 'Black Maul' | F | Cv GB | ♂ | X XXX X SSS | 1.5-2.1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 5-7 | --- | ♂ | |

| | | | | | | | | | | | | | | | | | | | |
|--|---|-------------|---|---------|---|-----|---|-----|---|---|---|------------------------------------|---|-----|---|---|-------|---|---|
| Salix triandra 'Champion' Almond-leaved Willow 'Champion' | M | Cv GB |  XXX X -- SSS | 1.8-2.7 | --- | --- | --- | --- | --- |  |  | NP Feb - - Mar - - Apr - 3 May 3 3 |  | --- |  |  | 5-9 | --- |  |
| Salix triandra 'Grisette Noire' Almond-leaved Willow 'Grisette Noire' | F | Cv F |  X XXX X SSS | 1.5-2.8 | --- | --- | --- | --- | --- | --- | --- | --- |  | --- |  |  | 5-7 | --- |  |
| Salix triandra 'Noire de Challans' Almond-leaved Willow 'Noire de Challans' | F | Cv F |  X XXX X SSS | 1.5-2.0 | --- | --- | --- | --- | --- | --- | --- | --- |  | --- |  |  | 5-7 | --- |  |
| Salix viminalis 'Cinnamomea' Common Osier 'Cinnamomea' | M | Cv D |  XX X --- SS | 1.6-2.1 | --- | --- | --- | --- | --- |  |  | NP Feb - - Mar - 3 Apr 2 2 May - - |  | --- | --- |  | 7-10 |  |  |
| Salix viminalis 'Heidenriet' Common Osier 'Heidenriet' | F | Lv CH |  XX X --- SSS | 1.7-2.3 | --- | --- | --- | --- | --- | --- | --- | --- |  | --- | --- |  | 7-10 |  |  |
| Salix viminalis 'Regalis' Common Osier 'Regalis' | F | Cv D |  XXX --- --- SSS | 2.1-3.0 | --- | --- |  | XX | --- | --- | --- | --- |  | --- | --- |  | 7-10 |  |  |
| Salix x americana American Willow | M | Cv USA/D |  --- XXX X SSS | 1.5-2.3 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |  | 3-4 | --- |  |
| Salix x aquatica 'Gigantea 56' Willow for Paper Production 56 | M | Cv DK | --- | 1.8-2.5 | --- | --- |  | XXX | --- | --- |  | NP Feb - - Mar - 3 Apr 3 3 May - - |  | --- | --- |  | 4-6 | --- |  |
| Salix x bögelsackii Bögelsack Willow | M | Cv D | --- | --- | --- | --- | --- | --- | --- | --- |  | NP Feb - - Mar - 4 Apr 4 4 May - - | --- | --- | --- |  | 5-6 | --- | --- |
| Salix x dasyclados Hybrid Willow 'Dasyclados' | F | Cv D/GB |  XX --- --- SS | 1.8-2.5 | --- | --- |  | XXX | --- | --- | --- | --- | --- | --- | --- |  | 4-6 |  |  |
| Salix x fragilis 'Bouton Plat' Hybrid Crack Willow 'Bouton Plat' | M | Cv F |  XX XX --- S | 1.5-1.8 | --- | --- | --- | --- |  | --- | --- | --- | --- | --- | --- |  | 10-15 |  |  |
| Salix x fragilis 'Flanders Red' Hybrid Crack Willow 'Flanders Red' | F | Cv B |  XX XX --- S | 1.5-2.2 |  | XXX | --- | --- |  | --- | --- | --- | --- | --- | --- |  | 10-15 |  |  |
| Salix x fragilis 'Jaune Hâtive' Hybrid Crack Willow 'Jaune Hâtive' | M | Cv F |  XX XX --- S | 1.5-1.8 | --- | --- | --- | --- |  | --- | --- | --- | --- | --- | --- |  | 10-15 |  |  |
| Salix x fragilis 'Opas Bindeweide' Hybrid Crack Willow 'Opas Bindeweide' | M | Lv CH |  XX XX --- S | 1.5-2.2 |  | XXX | --- | --- |  | --- | --- | --- | --- | --- | --- |  | 10-15 |  |  |
| Salix x mollissima 'Kottenheider Weide' Hybrid Willow 'Kottenheider Weide' | F | Cv D |  XXX X --- SS | 1.5-2.8 | --- | --- |  | X | --- | --- | --- | --- |  | --- | --- |  | 4-6 | --- |  |
| Salix x mollissima 'Trustworthy' Hybrid Willow 'Trustworthy' | M | Cv GB |  XX XX --- SS | 1.5-2.1 | --- | --- | --- | --- | --- | --- | --- | --- |  | --- | --- |  | 3-5 | --- |  |
| Salix x rubra 'Continental Osier' Green-leaved Willow 'Continental Osier' | F | Cv GB |  XX XX X SS | 1.6-2.5 |  | X | --- | --- | --- | --- | --- | --- | --- | --- | --- |  | 5-8 |  |  |
| Salix x rubra 'Ulbrich' Ulbrich Willow | F | Cv D |  XX XX --- SSS | 1.8-2.5 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |  | 5-8 |  |  |

| | | |
|---|---------------------------|---|
| Key to symbols: | M or F | male or female clone |
| | Cv or Lv | cultivar or local variety |
|  | XX XXX --- | for basketry, coarse medium fine suitable for white (S) (incl. value) |
| | SSS | annual growth [m] colour of fresh rods / colour of dried rods (depends on cultivation and storage conditions) |
|  | XXX | for tying in horticulture/ in viticulture (incl. value) |
|  | XXX | to grow thicker sticks or for biomass production (incl. value) |
|  | | for wood production / for living structure |
|  | NP Feb - - Mar - 4 | for beekeeping, NP = nectar and pollen, colour of the bee's pollen load beekeeping value (4 > 1) according to: Pritsch, Günter (2007): Bienenweide. 200 Trachtpflanzen erkennen und bewerten: Stuttgart: Kosmos. |
|  | | to provide fodder for ruminants (as a consequence, not resistant to deer) |
|  | ? | for drug production (suits the specifications of European Pharmacopoeia) / drug production in clarification |
|  | 5-6 | growth habit without pruning (tree or shrub); height [m] |
|  | | for pollarding / to grow rods for basketry (willow bed) |

25.09.2022/züm